

History of Araneology in Virginia

Barbara J. Abraham

Department of Biological Sciences
Hampton University
Hampton, Virginia 23668

ABSTRACT

At least from the 1600s to the present, spiders have been observed, collected, and studied in Virginia. This paper endeavors to outline the history of araneology in Virginia from its inception through the first decade of the 21st century, including researchers of spiders at Virginia institutions and those who have studied Virginia spiders.

Key words: araneology, history, spiders, Virginia.

INTRODUCTION

The study of spiders in Virginia has a venerable history, beginning in the 17th century and continuing to the present. Through the efforts of natural historians, entomologists, and ecologists, we know much about these important arachnids, but distributions and even the presence of some species in Virginia remain to be discovered. In the 20th and 21st centuries, quantitative, manipulative studies have increasingly replaced anecdotal observations, and spiders are used as model organisms to address ecological and evolutionary hypotheses.

17th Century

Perhaps the first person to collect spiders in Virginia was John Banister (1650-1692), an Englishman who arrived in Virginia in 1678. Banister was a careful observer; Lewis (1957) called science writers before Banister “historians rather than scientists.” She judged that natural history in America actually began with Banister.

By 1680, Banister had composed a catalog of insects called *Collectio insectorum*. Spiders appeared under “*De Insectis Pedibus Octonis*”; harvestmen were called spiders; silverfish were called “stingless scorpions.” As late as the time of Linnaeus, scorpions and spiders were placed with silverfish in the insect order Aptera (Ewan & Ewan, 1970).

Banister died when accidentally shot while botanizing on the Roanoke River in 1692, before seeing publication of his works on insects. He had been

planning a *Natural History of Virginia* (Lewis, 1957). Instead, “Some Observations concerning Insects made by Mr. John Banister in Virginia, A.D. 1680” was published by Petiver (Banister & Petiver, 1701). Banister was the first to systematically describe any of the spiders of North America; not until between 1791 and 1802 did John Abbot draw the spiders of Carolina and Georgia (Ewan & Ewan, 1970).

Lack of proper attribution by Martin Lister (a 17th century physician and natural historian), who received Banister’s specimens in 1680 and data for illustrations by Petiver, kept Banister’s Virginia specimens from inclusion in Linnaeus’s (1758-1759) *Systema Naturae* (Ewan & Ewan, 1992). Petiver’s illustrations were sources for Linnaeus in his establishment of binomials (Ewan & Ewan, 1970). Since spiders were very much a side line to Banister’s botanizing and Linnaeus did not knowingly use his collections, it is not surprising that Bonnet (1945) and others were not aware of Banister’s contribution.

18th Century

Benjamin H. Latrobe (1764-1820), an Englishman who immigrated to Virginia in 1795-96, was considered the first professional architect in the U.S. and designed the Capitol in Washington (Library of Congress, 2009). His published journal (Latrobe, 1905) is a good example of the nature writing of that time. In it, Latrobe wrote extensively about a spider wasp and a spider near Fredericksburg that he manipulated to see how they would behave. He also opened the mud cells of some wasps and recorded the colors and sizes of the

paralyzed spiders within (no mention is made of any names for the spiders).

19th Century

Between Banister and the early work of Nathan Banks (1868-1953), it is difficult to find records of any Virginians who studied spiders. Banks was a student of John H. Comstock at Cornell University. A New Yorker by birth, he lived in Falls Church, Virginia and published on local spiders (among other taxa) while employed at the USDA in Washington (1890-1916). During this time he published a revision of Marx's 1889 catalog of Nearctic spiders (Banks, 1910). Banks moved to the Museum of Comparative Zoology at Harvard University, where he continued to publish on spiders until at least 1930. He described a number of new species of spiders (Bonnet, 1945), but none of his titles on spiders were specific to Virginia. Exceptionally prolific, Banks authored over 440 technical papers from 1890-1951 and had eight children.

James Henry Emerton, who wrote widely on spiders, including a book on the common spiders of the United States (Emerton, 1902), published a paper on cave spiders, including some in Virginia (Emerton, 1875). In Fountain Cave (Augusta Co.), he found "one young spider allied to *Tegenaria*" and in Weyer's Cave (Rockingham Co.) a species of *Linyphia* with normal eyes; he noted that some specimens of the latter species had colors as bright as those of spiders of the same family from cellars or shady woods.

Increasingly in the 20th and 21st centuries, cave animals are endangered, and several more recent papers include spiders of Virginia caves. John R. Holsinger at Old Dominion University has published several papers on cave faunas, including spiders. A list of all identified spiders that have been found in caves in Virginia (obligate and accidental) can be found in Holsinger & Culver (1988). A new review paper will include additional obligate cave spider records (Holsinger et al., *in press*).

In 1888, when the Virginia State Agricultural Experiment Station was established by an Act of Congress, serious work on insects was initiated in Virginia (French, 1950), but because spiders are not crop pests, they have never been studied as much as insects.

20th Century to Present

Between November 1936 and March 1937, there was a flurry of notes in *Science* by various authors concerning the distribution of the black widow spider (*Latrodectus mactans*). D. C. Lowrie began the

conversation by saying that the black widow had not been officially recorded in eight states, including Virginia (Lowrie, 1936). G. W. Jeffers countered, "This must come as a surprise to the naturalists of the state," since the black widow was so prevalent that nobody bothered to mention it (Jeffers, 1936). B. J. Kaston returned that the black widow was first recorded from Hog Island in Virginia by J. H. Emerton in 1875 and was also recorded "as abundant in the Norfolk area by L. D. Anderson and H. G. Walker, and from various localities by C. R. Willey. Moreover, Dr. Bogen (Bogen, 1926) cites cases of arachnidism ..." (Kaston, 1937). In the same issue, H. A. Allard stated that he had found black widows in four places in Virginia within the last two years (Allard, 1937). R. L. Taylor, who was in the Department of Biology at the College of William and Mary, remarked sarcastically, "If by 'officially recorded,' publication in a scientific periodical is meant ..." and documented a June 1936 paper in the *Quarterly Review of Biology* (Taylor, 1937). Lowrie must not have seen this document before he submitted his note in November of that year. Taylor continued, "With respect to Virginia, it was rather surprising to find it said that there was no official record from this state since the black widow is common to abundant in most of the state" and cited papers on bites in the *Virginia Medical Monthly*. Additionally, he said, the 1934-1935 *Proceedings of the Virginia Academy of Science* included a paper read by Elizabeth Burger, a graduate student at William and Mary, which included the incidence of arachnidism in Virginia. Taylor continued, "One spring day, Miss Burger and the writer took 50 individuals [black widows] ... from the stones of a rock-banded curbing 35 x 2 feet in area" [in the vicinity of Williamsburg] (Taylor, 1937). Elizabeth Burger (Jackson)'s unpublished Master's thesis focused on the black widow spider (Jackson, 1935).

During the latter part of the 20th century until the present, a number of researchers, faculty, and graduate students in Virginia have studied Virginia spiders. However, many spider distributions in the state remain unknown.

Richard L. Hoffman (1927-2012), Curator Emeritus of Recent Invertebrates at the Virginia Museum of Natural History (VMNH) in Martinsville, was a native Virginian who devoted most of his life to the natural history of Virginia and the southern Appalachians, while being the leading authority on the world's millipeds. Hoffman's collection of spiders at the VMNH is the largest in the state, and he kept records of all Virginia spider species based on published literature and this collection. He published 485 scholarly books and papers and over 50 popular articles on millipeds,

amphibians, reptiles, worms, mollusks, arachnids, beetles, and true bugs; described more than 600 new taxa, including a species of *Hypochilus* from Virginia and West Virginia (Hoffman, 1963); and had nearly 50 taxa named in his honor (Roble & Mitchell, 2009). Among his numerous publications on spiders were lists of Virginia purse-web spiders (Atypidae), anyphaenids (Anyphaenidae), ground spiders (Gnaphosidae), and grass spiders (*Agelenopsis*) (e.g., Hoffman, 1992, 2000, 2002, 2009, 2010); removal of *Theridion montanum* (Theridiidae) from the Virginia faunal list (Hoffman, 1996a); a description of the type locality and distribution of *Xysticus emertoni* (Thomisidae) (Hoffman, 1996b); and addition of new species such as *Araneus saevus* (Araneidae), *Arachosia cubana* (Anyphaenidae), *Anahita punctulosa* (Ctenidae), and *Drassyllus rufulus* (Gnaphosidae) to the Virginia fauna (e.g., Hoffman, 1997, 2006; Hoffman et al., 2006; Roble & Hoffman, 2012).

Due to the difficulty in maintaining collections of spiders in alcohol, rather than dry, as for insects, some museums do not accept donations of private collections. Thanks to the interest of Richard Hoffman, the VMNH, established in 1988, houses thousands of spiders from Virginia. Soon after the Museum opened, Bill Shear, who was chairman of its Board of Directors, donated his personal spider collection of over 900 vials containing about 300 species (Hoffman, 1991). Shear's collection was mainly from outside Virginia, but Hoffman embarked upon an ambitious effort, in concert with the Virginia Division of Natural Heritage, to sample arachnids across the state. He expected to gather about 800 species of arachnids from Virginia. Thousands of mega-arachnids (not mites) were collected prior to 1991; these yielded a number of new state records and range extensions (Hoffman, 1991). Many more spiders were added to the VMNH collection up until Hoffman's death in June 2012.

At the Ninth International Congress of Arachnology in Panama in 1983, James E. Carico (1937-2009) of Lynchburg College proposed a "new" American spider family, the Trechaleidae (Carico, 1986b). He published revisions of genera in that family as well as in the Pisauridae (e.g., Carico, 1973, 1976, 1993, 2005a; Da Silva et al., 2008), studies on the behavior and anatomy of spiders in these families (e.g., Carico & Holt, 1964; Carico, 1986a; Bruce & Carico, 1988; Costa-Schmidt et al., 2008), and descriptions of new genera (e.g., Carico, 2005b). Carico served terms as President of the American Arachnological Society and Editor of its publication, *The Journal of Arachnology*.

James O. Howell completed his Master's thesis at Virginia Tech on "Spiders of alfalfa with notes on the biology of *Tetragnatha laboriosa* Hentz" (Howell,

1969). With his advisor Robert L. Pienkowski, he wrote the classic paper in the *Journal of Economic Entomology*, "Spider populations in alfalfa, with notes on spider prey and effect of harvest" (Howell & Pienkowski, 1971). The introduction in this paper is credited with showing that experimental studies of spiders in American agriculture were mostly absent at that time (Bell, 2005). The study itself used a D-Vac® to sample the litter and upper soil and 500-sweeps of a 15-in diameter net to take spiders on the alfalfa at one-week intervals during the warm months (monthly during the colder months) from March 1967 to June 1968. Several samples were taken at three-hour intervals during a twenty-four hour period to assess time of day on sampling efficiency. Temperature and humidity readings were taken. Over 4,000 spiders in 15 families and 124 species were collected. The attention to environment, season, and time of day effects, scope of the sampling, and addition of laboratory feeding tests to address what the spiders were actually eating in the alfalfa mark this paper as one of the most thorough of the early studies of spiders in crop fields.

In an example of the observational approach to studying spiders, Laura Elsa Sabath collected an individual *Gea heptagon* from her web in a lawn in Portsmouth, Virginia, put her into a terrarium, and observed the spider until she died (Sabath, 1969). Sabath's motivation was to study a spider that was not currently being used by other experimenters, and one that, although widespread, had not been studied extensively. She made observations on rapid change of the spider's color after it fell to the ground when disturbed, web spinning, feeding, the egg sac, and spiderlings.

J. P. McCaffrey and R. L. Horsburgh of Virginia Tech's Shenandoah Valley Research Station studied spiders in Virginia apple orchards and used limb beating as a sampling method for collecting spiders on apple trees. They documented that the proportions of hunting spiders in current vs. abandoned orchards were similar in most cases. The spider species exhibited niche separation on different parts of the trees; the authors thought that this might facilitate suppression of insect pests (McCaffrey & Horsburgh, 1980). Another study evaluated time of day and season as factors in sampling results using limb beating. *Clubiona* spp., which are nocturnal, were captured significantly more often at 0300 h, while salticids and total spiders were collected more during 0900 and 1800 h. Limb beating was deemed a satisfactory method for collecting both web-building and hunting spiders, but spiderlings of some (*Philodromus*) were not captured efficiently by this method (McCaffrey et al., 1984).

C. L. Steitenroth (with Norm Horner at Midwestern

State University) published an account of 18 genera and 30 species of jumping spiders of the Virginia Peninsula (Steitenroth & Horner, 1987). The study produced records of five species new to the state.

William A. Shear of Hampden-Sydney College has published more than 200 papers and book chapters, primarily relating to classification and evolution of arthropods, and is the co-author or editor of two books, including *Spiders: Webs, Behavior and Evolution* (Shear, 1986). His paleontological studies began in 1980; more recently he returned to taxonomy and to a new direction in ecological chemistry. He has published notes in *Banisteria* on a leptonetid and a pholcid new to Virginia (Shear, 2007; Clark & Shear, 2010) and was chief scientific consultant, script editor, and on-camera participant for the film, "Ultimate Guide: Spiders," broadcast on 9 July 2001 on the Discovery Channel.

A paper published by Shear and colleagues on a fossil spider from New York extended the fossil record of non-araneomorph spiders back to the Devonian (Selden et al., 1991). A later paper including a fossil Virginia spider from the Triassic was published by Nicholas C. Fraser of the VMNH and co-workers (Selden et al., 1999). These specimens represented the oldest known fossil araneomorph spiders and extended the fossil record of that group back significantly. Shear and colleagues also published papers on fossil evidence of the origin of spider spinnerets, which are considered the defining adaptation of spiders (Shear, 1989; Selden et al., 2008). In the 2008 study, fossil evidence was combined with developmental genetic studies to clarify how use of silk may have evolved in spiders.

Brent Opell's research at Virginia Tech is an example of the modern experimental, quantitative approach to studying spiders. Early studies of Virginia *Hyptiotes* (Uloboridae), for example, showed that web production of early instars is independent of container size (Opell, 1982). Opell also has published a revision and a checklist of American Uloboridae (Opell, 1979; 1983). His research currently focuses on spider systematics and integrative biology using morphological and molecular characters and phylogeography (Opell Lab, 2009). These studies have produced numerous papers on the viscous capture threads of orb weavers (e.g., Opell & Hendricks, 2009). These and other studies on spider morphology, behavior, genetics, and evolution (e.g., Opell, 1984, 1990, 1998, 2010; Opell & Bond 2001; Opell et al., 2007) are of broad interest to spider systematists and ecologists.

Michael W. Beck and Edward F. Connor of the University of Virginia and Blandy Experimental Farm conducted a study examining the importance of covariance in traits related to foraging, between developmental stages of the crab spider *Misumenoides*

formosipes (Beck & Connor, 1992). They found that prereproductive weight and fecundity were highly correlated to carapace width. Growth of spiders fed *ad libitum* in the laboratory was unrelated to size, which suggested that size in the field was related to prey-capture success. Small females (measured by carapace width) did not reproduce, although they constituted a significant proportion of the population. The authors emphasized examining stage-specific constraints in order to understand the effects of foraging on reproductive success.

While at Virginia Commonwealth University, Stephen R. Johnson studied spiders of early successional stages of a barrier island (Hog Island) that is part of the Virginia Coast Reserve. He found that spider diversity and density differed more between sites than between shrub species (*Baccharis halmifolia* and *Myrica cerifera*) (Johnson, 1996).

Steven M. Roble of the Virginia Department of Conservation and Recreation's Division of Natural Heritage recorded a range extension to Virginia for *Gasteracantha cancriformis* (Roble, 1994). Roble often collaborated with Richard Hoffman, co-authoring several papers on spiders (and other taxa) new to the Virginia fauna (Hoffman & Roble, 2012 [three wolf spiders]; Roble & Hoffman, 2012 [*Drassyllus rufulus*]).

The author of this paper published a preliminary list of the spiders of the Great Dismal Swamp (Abraham, 2000), as well as a list of the spiders found during the Potomac Gorge Bioblitz (Abraham, 2008). Although other faunal lists for the Great Dismal Swamp date from much earlier, Abraham's list is the first for the spiders of the Swamp. The list includes 14 families, 43 genera, and 56 species. Likewise, the list of Potomac Gorge spiders, albeit incomplete, is the first specific to that area (although many spiders had previously been collected in and around the Washington, D.C. area). The Potomac Gorge Bioblitz list included 37 species in 29 genera and 12 families.

Anne Danielson-Francois, Christine A. Fetterer, and Peter D. Smallwood (the latter two authors are from the University of Richmond) published a paper on body condition and mate choice in *Tetragnatha elongata* (Danielson-Francois et al., 2002). They found that males preferred longer, heavier females with higher body condition. This study may have been the first to show the influence of body condition on mate choice in spiders. Smallwood had previously published a study on web site tenancy in this species that challenged a model of risk-sensitive foraging and emphasized the importance of considering multiple scales in ecology (Smallwood, 1993).

Rachel E. Mallis and Lawrence E. Hurd at Washington and Lee University studied ground-

dwelling spider assemblages in the Science Park at their institution with pitfall traps (Mallis & Hurd, 2005). They collected 50 species from six sites, but found no correlation between spider assemblages and successional stage of the habitat. They thought that spider community composition is generally unpredictable, due to stochastic colonization and specific resource requirements following immigration.

While at the University of Virginia, Lily Ahrens and Johanna M. Kraus (Kraus is now at Washington and Lee University) reported on wolf spider movements along a pond edge at Mountain Lake Biological Station, near Pembroke, Virginia using mark-recapture. In contrast to previous studies, in this study wolf spiders (*Pardosa* spp.) moved little in time or space, although the short-term (summer) and small spatial scale (meters) of this study may have had an impact (Ahrens & Kraus, 2006).

At the 2008 Virginia Academy of Science meetings, Marc A. Milne, then a graduate student of Deborah Waller at Old Dominion University, presented on nectar feeding by spiders of three families: Linyphiidae, Lycosidae, and Agelenidae. His dissertation concerned spiders associated with the purple pitcher plant (*Sarracenia purpurea*) in Virginia and North Carolina (Milne, 2010). From this research he published a paper on the purple pitcher plant as a spider oviposition site (Milne, 2012).

Already in 1990, the lack of support for taxonomic arachnology was lamented (Coddington et al., 1990). The situation is not as dire for spiders as for other arachnid groups, but few young scientists enter the field due to a lack of open, funded positions (Coddington et al., 1990). There is yet much remaining to be discovered about spiders in Virginia, as well as the rest of the world, in 2013, and conservation of these important predators is not well served by the lack of information. Virginia is lucky to have the VMNH, the Division of Natural Heritage, and the extant individuals mentioned above to carry on the study of the state's spiders, but new recruits and funding are necessary if complete information on the presence and distribution of spiders in Virginia is to be obtained.

ACKNOWLEDGEMENTS

Thanks to Dr. Alfred Willis, Science Librarian at the William R. and Norma B. Harvey Library, Hampton University, who enthusiastically helped me locate some early sources. I also thank Tom McAvoy and Steve Roble for their significant editorial assistance with changing my symposium presentation into a paper.

LITERATURE CITED

- Abraham, B. J. 2000. A preliminary list of the spiders of the Great Dismal Swamp. Pp. 139-141 *In* R. K. Rose (ed.), *The Natural History of the Great Dismal Swamp*. Omni Press, Madison, WI.
- Abraham, B. J. 2008. Arachnids. Pp. 37-38 *In* A. V. Evans and survey team leaders. *The 2006 Potomac Gorge Bioblitz*. *Banisteria* 32.
- Ahrens, L., & J. M. Kraus. 2006. Wolf spider (Araneae: Lycosidae) movement along a pond edge. *Journal of Arachnology* 34: 532-539.
- Allard, H. A. 1937. The black widow spider in Virginia. *Science* 85: 74-75.
- Banister, J., & J. Petiver. 1701. Some observations concerning insects made by Mr. John Banister in Virginia, A.D. 1680, with remarks on them by Mr. James Petiver, Apothecary and Fellow of the Royal Society. *Philosophical Transactions* 1700-1701(22): 807-814. doi: 10.1098/rstl.1700.0078.
- Banks, N. 1910. Catalogue of the Nearctic spiders. *Bulletin of the U. S. National Museum* 72. Washington, DC. 80 pp.
- Beck, M. W., & E. F. Connor. 1992. Factors affecting the reproductive success of the crab spider *Misumenoides formosipes*: the covariance between juvenile and adult traits. *Oecologia* 92: 287-295.
- Bell, J. R. 2005. The emergence of manipulative experiments in ecological spider research (1684-1973). *Journal of Arachnology* 33: 826-849.
- Bogen, E. 1926. Arachnidism: spider poisoning. *Archives of Internal Medicine* 38: 623-632.
- Bonnet, P. 1945. *Bibliographia araneorum*, Vol. 1. Pub. by author at Toulouse; reprinted 1968 Noble Offset Printers, Inc., USA. 832 pp.
- Bruce, J. A., & J. E. Carico. 1988. Silk use during mating in *Pisaurina mira* (Walckenaer) (Araneae, Pisauridae). *Journal of Arachnology* 16: 1-4.
- Carico, J. E. 1973. The Nearctic species of the genus *Dolomedes* (Araneae: Pisauridae). *Bulletin of the Museum of Comparative Zoology* 144: 435-488.

- Carico, J. E. 1976. The spider genus *Tinus* (Pisauridae). *Psyche* 83: 63-78.
- Carico, J. E. 1986a. Web removal patterns in orb-weaving spiders. Pp. 306-318 *In* W. A. Shear (ed.), *Spiders: Webs, Behavior and Evolution*. Stanford University Press, Stanford, CA.
- Carico, J. E. 1986b. Trechaleidae: A "new" American spider family. (Abstract). P. 305 *In* W. G. Eberhard, Y. D. Lubin, & B. C. Robinson (eds.), *Proceedings of the Ninth International Congress of Arachnology*, Panama, 1983. Smithsonian Institution Press, Washington, DC.
- Carico, J. E. 1993. Revision of the genus *Trechalea* Thorell (Araneae, Trechaleidae) with a review of the taxonomy of the Trechaleidae and Pisauridae of the Western Hemisphere. *Journal of Arachnology* 21: 226-257.
- Carico, J. E. 2005a. Revision of the spider genus *Hesydrus* (Araneae, Lycosoidea, Trechaleidae). *Journal of Arachnology* 33: 785-796.
- Carico, J. E. 2005b. Descriptions of two new spider genera of Trechaleidae (Araneae, Lycosoidea) from South America. *Journal of Arachnology* 33: 797-812.
- Clark, T., & W. A. Shear. 2010. The tailed cellar spider, *Crossopriza lyoni* (Blackwall 1867), new to Virginia (Araneae: Pholcidae). *Banisteria* 36: 45.
- Coddington, J. A., S. F. Larcher, & J. C. Cokendolpher. 1990. The systematic status of Arachnida, exclusive of Acari, in North America north of Mexico. Pp. 5-20 *In* M. Kosztarab & C. W. Schaefer (eds.), *Systematics of the North American Insects and Arachnids: Status and Needs*. Virginia Agricultural Experiment Station Information Series 90-1. Virginia Polytechnic Institute and State University, Blacksburg.
- Danielson-Francois, A., C. Fetterer, & P. D. Smallwood. 2002. Body condition and mate choice in *Tetragnatha elongata*. *Journal of Arachnology* 30: 20-30.
- Emerton, J. H. 1875. Notes on spiders from caves in Kentucky, Virginia and Indiana. *American Naturalist* 9: 278-281.
- Emerton, J. H. 1902. *The Common Spiders of the United States*. The Athenaeum Press, Ginn & Co., Publ., Boston. 225 pp.
- Ewan, J., & N. Ewan. 1970. *John Banister and his Natural History of Virginia 1678 – 1692*. University of Illinois Press, Urbana. 485 pp.
- Ewan, J., & N. Ewan. 1992. John Banister, Virginia's first naturalist. *Banisteria* 1: 3-5.
- French, G. T. 1950. Entomology. Pp. 107-128 *In* *The James River Basin: Past, present and future*. James River Project Committee of the Virginia Academy of Science. Virginia Academy of Science, Richmond, VA. 843 pp.
- Hoffman, R. L. 1963. A second species of the spider genus *Hypochilus* in eastern North America. *American Museum Novitates* 2148: 1-8.
- Hoffman, R. L. 1991. Arachnology at the Virginia Museum of Natural History. *American Arachnology* 44: 2-3.
- Hoffman, R. L. 1992. Purse-web spiders (Atypidae) in Virginia. *Banisteria* 1: 5-7.
- Hoffman, R. L. 1996a. Deletion of the spider *Theridion montanum* from the Virginia faunal list (Araneida: Theridiidae). *Banisteria* 8: 55-56.
- Hoffman, R. L. 1996b. Type locality and distribution of the crab spider *Xysticus emertoni* Keyserling (Araneida: Thomisidae). *Banisteria* 8: 53-54.
- Hoffman, R. L. 1997. First records of a Holarctic orb-weaving spider (*Araneus saevus* [L. Koch]) in Virginia. *Banisteria* 10: 30.
- Hoffman, R. L. 2000. Grass spiders of the genus *Agelenopsis* in Virginia (Araneida: Agelenidae). *Banisteria* 15: 36-42.
- Hoffman, R. L. 2002. Spiders of the family Anyphaenidae in Virginia (Arachnida: Araneida). *Banisteria* 19: 12-16.
- Hoffman, R. L. 2006. A note on the occurrence of the spider *Arachosia cubana* (Banks) in Virginia (Araneae: Anyphaenidae). *Banisteria* 27: 45.
- Hoffman, R. L. 2009. Virginia ground spiders: A first list (Araneae: Gnaphosidae). *Banisteria* 33: 18-29.
- Hoffman, R. L. 2010. Purse-web spiders, genus *Sphodros*, in Virginia (Mygalomorphae: Atypidae). *Banisteria* 36: 31-38.

- Hoffman, R. L., W. J. Arnold, & R. A. Bradley. 2006. *Anahita punctulata* (Hentz): A species, genus, and family of spiders new to the fauna of the Virginias and Ohio (Arachnida: Araneae: Ctenidae). *Banisteria* 27: 46-47.
- Hoffman, R. L., & S. M. Roble. 2012. Three wolf spiders new to the Virginia fauna (Araneae: Lycosidae). *Banisteria* 40: 54-57.
- Holsinger, J. R., & D. C. Culver. 1988. The invertebrate cave fauna of Virginia and a part of east Tennessee: zoogeography and ecology. *Brimleyana* 14: 1-162.
- Holsinger, J. R., D. C. Culver, D. A. Hubbard, Jr., W. Orndorff, & C. Hobson. The invertebrate cave fauna of Virginia. *Banisteria* 42: *in press*.
- Howell, J. O. 1969. Spiders of alfalfa with notes on the biology of *Tetragnatha laboriosa* Hentz. Master's Thesis, Virginia Polytechnic Institute and State University, Blacksburg, VA. 114 pp.
- Howell, J. O., & R. L. Pienkowski. 1971. Spider populations in alfalfa, with notes on spider prey and effect of harvest. *Journal of Economic Entomology* 64: 163-168.
- Jackson, E. B. 1935. A study of the "black widow" spider *Latrodectus mactans*, fabricius (sic.). Master's Thesis, The College of William and Mary, Williamsburg, VA.
- Jeffers, G. W. 1936. The black widow spider in Virginia. *Science* 84: 533-534.
- Johnson, S. R. 1996. Spiders associated with early successional stages on a Virginia barrier island. *Journal of Arachnology* 24: 135-140.
- Kaston, B. J. 1937. The distribution of black widow spiders. *Science* 85: 74.
- Latrobe, B. H. 1905. *The Journal of Latrobe, Being the Notes and Sketches of an Architect, Naturalist, and Traveler in the United States from 1796 to 1820*. D. Appleton and Co., New York. Available online at: <http://archive.org/details/journaloflatrobe00latrrih>
- Lewis, I. F. 1957. Seventeenth century science in Old Virginia. *Virginia Journal of Science* 8: 35-41.
- Library of Congress. 2009. American memory: Built in America. Online at: http://memory.loc.gov/cgi-bin/query/D?hh:1:/temp/~ammem_mnK0::. Accessed 13 June 2013.
- Linnaeus, C. 1758-1959. *Caroli Linnaei Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum charateribus, differentiis, synonymis, locis*. ed. 10.
- Lowrie, D. C. 1936. New localities for the black widow spider. *Science* 84: 437.
- Mallis, R. E., & L. E. Hurd. 2005. Diversity among ground-dwelling spider assemblages: habitat generalists and specialists. *Journal of Arachnology* 33: 101-109.
- McCaffrey, J. P., & R. L. Horsburgh. 1980. The spider fauna of apple trees in central Virginia. *Environmental Entomology* 9: 247-252.
- McCaffrey, J. P., M. P. Parrella, & R. L. Horsburgh. 1984. Evaluation of the limb-beating sampling method for estimating spider (Araneae) populations on apple trees. *Journal of Arachnology* 11: 363-368.
- Milne, M. A. 2010. Ecological interactions between spiders and the purple pitcher plant, *Sarracenia purpurea*. Ph.D. Dissertation, Old Dominion University, Norfolk, VA. 140 pp.
- Milne, M. A. 2012. The purple pitcher plant as a spider oviposition site. *Southeastern Naturalist* 11: 567-574.
- Opell, B. D. 1979. Revision of the genera and tropical American species of the spider family Uloboridae. *Bulletin of the Museum of Comparative Zoology* 148: 433-549.
- Opell, B. D. 1982. Post-hatching development and web production of *Hyptiotes cavatus* (Hentz) (Araneae, Uloboridae). *Journal of Arachnology* 10: 185-191.
- Opell, B. D. 1983. Checklist of American Uloboridae (Arachnida: Araneae). *Great Lakes Entomologist* 16: 61-66.
- Opell, B. D. 1984. Eggsac differences in the spider family Uloboridae (Arachnida: Araneae). *Transactions of the American Microscopical Society* 103: 122-129.
- Opell, B. D. 1990. Material investment and prey capture potential of reduced spider webs. *Behavioral Ecology and Sociobiology* 26: 375-381.
- Opell, B. D. 1998. Economics of spider orb-webs: the

- benefits of producing adhesive capture thread and of recycling silk. *Functional Ecology* 12: 613-624.
- Opell, B. D. 2010. Bergmann's size cline in New Zealand marine spray zone spiders (Araneae: Anyphaenidae: *Amaurobioides*). *Biological Journal of the Linnaean Society* 101: 78-92.
- Opell, B. D., A. Berger, S. Bous, & M. Manning. 2007. Genetic relationships of *Amaurobioides* (Anyphaenidae) spiders from the southeastern coast of New Zealand. *Zootaxa* 1425: 1-10.
- Opell, B. D., & J. E. Bond. 2001. Changes in the mechanical properties of capture threads and the evolution of modern orb-weaving spiders. *Evolutionary Ecology* 3: 567-581.
- Opell, B. D., & M. L. Hendricks. 2009. The adhesive delivery system of viscous capture threads spun by orb-weaving spiders. *Journal of Experimental Biology* 212: 3026-3034.
- Opell Lab, The. 2009. [Brent Opell's Research Lab]. Spider Systematics and Integrative Biology. Online at: <http://www.faculty.biol.vt.edu/opell/>. Accessed 13 June 2013.
- Roble, S. M. 1994. *Gasteracantha cancriformis* (L.), a spectacular spider new to the fauna of Virginia (Araneae: Araneidae). *Banisteria* 3: 20.
- Roble, S. M., & J. C. Mitchell (eds.). 2009. A Lifetime of Contributions to Myriapodology and the Natural History of Virginia: A Festschrift in Honor of Richard L. Hoffman's 80th Birthday. Virginia Museum of Natural History Special Publication Number 16, Martinsville, VA. 458 pp.
- Roble, S. M., & R. L. Hoffman. 2012. *Drassyllus rufulus* (Banks), an addition to the ground spider fauna of Virginia (Araneae: Gnaphosidae). *Banisteria* 40: 68.
- Sabath, L. A. 1969. Color change and life history observations of the spider *Gea heptagon* (Araneae: Araneidae). *Psyche* 76: 367-374.
- Selden, P. A., J. M. Anderson, H. M. Anderson, & N. C. Fraser. 1999. Fossil Araneomorph spiders from the Triassic of South Africa and Virginia. *Journal of Arachnology* 27: 401-414.
- Selden, P.A., W. A. Shear, & P. M. Bonamo. 1991. A spider and other arachnids from the Devonian of New York, and reinterpretations of Devonian Araneae. *Paleontology* 34: 241-281.
- Selden, P. A., W. A. Shear, & M. D. Sutton. 2008. Fossil evidence for the origin of spider spinnerets, and a proposed arachnid order. *Proceedings of the National Academy of Sciences (USA)* 105: 20781-20785.
- Shear, W. A. (ed.). 1986. *Spiders: Webs, Behavior and Evolution*. Stanford University Press, Stanford, CA. 492 pp.
- Shear, W. A. 2007. *Appaleptoneta coma* (Barrows), a spider new to Virginia (Araneae, Leptonetidae). *Banisteria* 30: 40-41.
- Shear, W. A., J. M. Palmer, J. A. Coddington, & P. M. Bonamo. 1989. A Devonian spinneret: Early evidence of spiders and silk use. *Science* 246: 479-481.
- Smallwood, P. D. 1993. Web-site tenure in the long-jawed spider: Is it risk-sensitive foraging, or conspecific interactions? *Ecology* 74: 1826-1835.
- Steitenroth, C. L., & N. V. Horner. 1987. The jumping spiders (Araneae: Salticidae) of the Virginia Peninsula. *Entomological News* 98: 235-245.
- Taylor, R. L. 1937. The black widow spider in Virginia. *Science* 85: 263-264.